Abstract Submitted to the International Conference on Strongly Correlated Electron Systems University of Michigan, Ann Arbor August 6-10, 2001

Acoustical Measurements of Heavy Fermion Compounds in Pulsed Magnetic Fields *

Alexei Souslov¹, Debashsish Dasgupta¹, Jeff R. Feller¹, Marcelo Jaime², Fedor Balakirev², David G. Hinks³, Albert Migliori², Alex Lacerda², John B. Ketterson⁴, Bimal K. Sarma¹

- ¹ Physics Department, University of Wisconsin-Milwaukee, Milwaukee, WI 53201, USA
- ² National High Magnetic Field Laboratory, Los Alamos, NM 87545, USA
- Materials Science & Technology, Argonne National Laboratory, Argonne, IL 60439, USA
- ⁴ Northwestern University, Evanston, IL 60208, USA

Ultrasonic properties of UPt_3 and URu_2Si_2 crystals have been measured at 2-4 K in pulsed (25ms) magnetic fields up to 50T. In the frequency range from 20 to 400MHz, both the attenuation and velocity of sound, have changed in the vicinity of metamagnetic transition in these materials (around 20T and 40T respectively). Data shows that three transitions occur in URu_2Si_2 in agreement with magnetization results of other authors. For these measurements an original fast data acquisition ultrasonic spectrometer has been developed and assembled for these studies in pulsed magnetic field.

^{*}The work was supported by NSF grant DMR-9704020 and DMR-9971123. The NHMFL is supported by the NSF, the State of Florida and the US Department of Energy.